



As we age, especially due to oxidative stress and other external factors, the efficiency of the immune response tends to progressively decrease. This weakening can lead to disproportionate immune reactions, resulting in exaggerated inflammatory responses that, instead of being beneficial, contribute to tissue deterioration.

The compounds present in the formula have the ability to reach the intercellular space through direct bloodstream circulation, especially in the case of orally administered products. During acute inflammation, enzymes play a crucial role by contributing to the degradation of plasma proteins in the interstitial space.

This action facilitates the removal of these proteins through the lymphatic or bloodstream system.

In summary, the aging process, along with oxidative stress, can negatively affect the immune response, leading to harmful inflammatory responses. However, the compounds in the formula, by accessing the intercellular space through bloodstream circulation and with the help of enzymes, play an important role in mitigating acute inflammation by facilitating the removal of plasma proteins, thereby contributing to tissue health maintenance.

Benefits

- Reduces and prevents excessive inflammation
- Improves reduction of clinical and subclinical inflammation
- Provides support in treating BPH (Benign Prostatic Hyperplasia)
- Supports degradation and removal of Fibrin Sheath

Highlights

- Includes cellular extracts and antioxidants for immune support.
- Contains enzymes to degrade proteins during inflammation.
- Provides calcium, zinc, selenium, and vitamins D, C, and E.
- Additional components with potential anti-inflammatory properties and bone benefits.

Recommended Use

Preventive Action and Aging: One tablet in the morning and 1 in the evening, for at least six months.

Adjunct Action (Subclinical Inflammation, Post-surgical Processes): Two tablets in the morning and 2 in the evening, for at least six months.

Osteoarticular Inflammatory Processes and Benign Prostatic Hyperplasia: Two tablets in the morning and 2 in the evening, for at least three months.